



## Michigan Quality Improvement Consortium Guideline

*Adults with Systolic Heart Failure*

**The following guideline recommends diagnostic evaluation, pharmacologic treatment and education that support effective patient self-management.**

Eligible Population	Key Components	Recommendation and Level of Evidence
Adults with suspicion of left-ventricular systolic dysfunction, including heart failure	Evaluation	<p><u>Initial assessment should include:</u></p> <ul style="list-style-type: none"> <li>♦ Thorough history and physical examination [C], including depression screening, and assessment for coronary artery disease and risk factors</li> <li>♦ Testing includes: chest X-ray, 12-lead electrocardiogram, lipid profile, CBC, electrolytes, calcium, magnesium, BUN, creatinine, blood glucose, liver function tests, TSH, urinalysis, and echocardiography with Doppler [C]</li> <li>♦ BNP is useful for diagnosis, and not for serial monitoring</li> <li>♦ Serial monitoring should include: weight, volume status, electrolytes, renal function and activity tolerance</li> </ul>
Adults diagnosed with left-ventricular systolic dysfunction, including heart failure	Management	<p><u>Recommended for routine use:</u></p> <ul style="list-style-type: none"> <li>♦ ACE inhibitors in all patients, unless contraindicated<sup>1</sup> [A]</li> <li>♦ Beta-blockade using carvedilol, sustained-release metoprolol, bisoprolol in all stable patients, unless contraindicated<sup>1,2</sup> [A]</li> <li>♦ Consider referral for evaluation for implantable defibrillator, ventricular assist device or transplant in patients with LVEF&lt;35%, NYHA Class II-IV patients and those with worsening CHF</li> <li>♦ Consider referral for biventricular pacemaker for patients with symptomatic heart failure and QRS duration ≥ 120 msec</li> <li>♦ Consider referral of complex patients to an advanced heart failure management program</li> <li>♦ Vaccination against influenza and pneumococcal</li> </ul> <p><u>Recommended for use in select patients:</u></p> <ul style="list-style-type: none"> <li>♦ Diuretics and sodium restriction for evidence of fluid retention [A]</li> <li>♦ Spironolactone for patients with symptoms of heart failure, preserved renal function (creatinine &lt; 2.0 in women; creatinine &lt; 2.5 in men) and normal serum potassium concentration [A]</li> <li>♦ Use ARBs in patients who cannot tolerate ACE inhibitors due to cough or angioedema [A]</li> <li>♦ Consider hydralazine and isosorbide dinitrate for patients who cannot tolerate ACE inhibitors or ARBs, or African-American patients who remain symptomatic despite therapy [A]</li> <li>♦ Digoxin should only be used for patients who remain symptomatic despite diuretics, ACE inhibitors and beta blockers [A]</li> </ul>
	Counseling and care management	<p><u>Engage patients in office-based care management and self-management:</u></p> <ul style="list-style-type: none"> <li>♦ Careful review of medication regimen with patient and caregivers at hospitalization or other changes in treatment</li> <li>♦ Daily self-monitoring of weight and adherence to recommended patient action plan</li> <li>♦ Recognition of symptoms and when to seek medical attention</li> <li>♦ Moderate dietary sodium restriction (e.g., 2,000-2,500 mg sodium/day)</li> <li>♦ Risk factor modification (regular exercise 5 times per week as tolerated [B]; smoking cessation; control of BP, DM, lipids)</li> <li>♦ Avoid excessive alcohol intake, illicit drug use, and the use of NSAIDS</li> <li>♦ Discuss goals of care, prognosis, advance directives, and palliative care</li> </ul>

<sup>1</sup> Contraindications include: life-threatening adverse reactions (angioedema or anuric renal failure), pregnancy, hypotensive patients at immediate risk of cardiogenic shock, systolic blood pressure < 80 mm Hg, serum creatinine > 3 mg/dL, bilateral renal artery stenosis, or serum potassium > 5.5 mmol/L.

<sup>2</sup> Contraindications include: patients with current or recent fluid retention history, unstable or poorly controlled reactive airway disease, symptomatic bradycardia or advanced heart block (unless treated with a pacemaker), or recent treatment with an intravenous positive inotropic agent.

**Levels of Evidence for the most significant recommendations:** A = randomized controlled trials; B = controlled trials, no randomization; C = observational studies; D = opinion of expert panel

This guideline lists core management steps. It is based on the 2013 ACCF/AHA Guideline for the Management of Heart Failure: A Report of the American College of Cardiology Foundation/American Heart Association Task Force on Practice Guidelines. Developed in Collaboration With the American College of Chest Physicians, Heart Rhythm Society and International Society for Heart and Lung Transplantation. Individual patient considerations and advances in medical science may supersede or modify these recommendations.